


CLAIMS:


1. A cross-linkable compound containing one or more fluorendiyl groups of the formula:



where R, independently each occurrence, is an inert substituent, a monovalent crosslink forming group, X, or a polyvalent crosslink forming group, X', with the proviso that in at least one repeat unit per molecule, at least one R is X or X'.

2. A compound according to claim 1 wherein R in at least one occurrence is a moiety containing a double bond, a triple bond, a precursor capable of in situ formation of a double bond, or a heterocyclic, addition polymerizable group.
3. A compound according to claim 1 wherein R in at least one occurrence is selected from the group consisting of benzocyclobutanyl groups and substituted C<sub>6-12</sub> arylene groups containing one or more substituents selected from the group consisting of benzocyclobutane, azide, oxirane, di(hydrocarbyl)amino, cyanate ester, hydroxy, glycidyl ether, C<sub>1-4</sub> alkylacrylate, C<sub>1-4</sub> alkylmethacrylate, alkenyl, alkenyloxy, alkynyl, maleimide, nadimide, tri(C<sub>1-4</sub>)-alkylsiloxyl, tri(C<sub>1-4</sub>)alkylsilyl, and halogenated derivatives thereof.
4. A compound according to claim 1 wherein R in at least one occurrence is benzo-3,4-cyclobutan-1-yl or p-vinylbenzyl.

5. A cross-linkable composition comprising oligomers or polymers having empirical formula Ia:



where R, independently each occurrence, is an inert substituent, a monovalent crosslink forming group, X, or a polyvalent crosslink forming group, X', with the proviso that in at least one repeat unit per molecule, at least one R is X or X';


Z is a divalent remnant of a comonomer or a monovalent chain terminating group; and


x is a number from 1 to 10,000 and z is a number from 0 to 10,000 signifying the average number of repeat units in the composition.

6. A composition according to claim 5 wherein Z each occurrence is halo, cyano, triflate, azide,  $-B(OR^1)_2$ , or  $-B \begin{array}{c} \diagup O \diagdown \\ \diagdown O \diagup \end{array} R^2$ .

wherein R<sup>1</sup>, independently in each occurrence, is hydrogen or a C<sub>1-10</sub> alkyl group, and R<sup>2</sup>, independently each occurrence, is a C<sub>2-10</sub> alkylene group.


5            7.        A crosslinkable oligomer or polymer of the formula:





where R, x and z are as previously defined in claim 5 and Z'' is as defined in claim 7.

9. A cross-linked polymer corresponding to formula:




R' independently each occurrence is R or a crosslinked derivative of X or X' with the proviso, that in at least one occurrence, R' is a crosslinked derivative of X or X',

X, X', R, and x are as previously defined in claim 1, and

Z', Z'' and z are as defined in claim 7.

10. A process for preparing oligomers or polymers of claim 7, which process comprises heating one or more compounds containing one or more fluorendiyl groups of the formula:



where R, independently each occurrence, is an inert substituent, a monovalent crosslink forming group, X, or a polyvalent crosslink forming group, X', with the proviso that in at least one repeat unit per molecule, at least one R is X or X' or a composition comprising the same, optionally in the presence of a noninterfering compound, under reaction conditions sufficient to form an oligomer or polymer of Claim 7.

11. A film comprising one or more of the oligomers or polymers according to any one of claims 7 to 9 or preparable according to claim 10.

12. An electronic device comprising one or more layers of polymer films, at least one of which comprises a film according to claim 11.